



Barrington Office
Suite 312
(22285 N Pepper Rd, Ste 312,
Lake Barrington IL, 60010)

Good Shepherd
West Pavilion
(450 IL-22, Barrington, IL
60010)

Northwestern Huntley
Main Entrance
(10400 Haligus Rd, Huntley, IL
60142)

Northwestern McHenry
Main Entrance
(4201 Medical Centre Dr, McHenry, IL
60050)

Physician: _____ Date: _____ Arrival Time: _____ Procedure Time: _____

COLONOSCOPY USING CLENPIQ PREP: PLEASE, ONLY FOLLOW THESE DIRECTIONS-NOT WHAT IS ON THE BOX

A colonoscopy is a procedure that examines the large intestine with a flexible fiberoptic instrument. The doctor is looking for any abnormalities in the colon and may take biopsies. This procedure takes about 20 to 30 minutes. With intake and recovery however, you must plan to be at the facility for at least 1.5 hours (Barrington) up to 3 hours (hospital).

IT IS ESSENTIAL THAT...

1. You bring your ID, insurance card and a list of medications (including dosages) to your appointment.
2. You will not be allowed to drive yourself home after the procedure due to the sedation. Therefore, you will need an adult to drive you home. You may **NOT** use public transportation. **If you do not have a ride, your procedure will be cancelled.**

***STOP THESE MEDICATIONS 14 DAYS BEFORE YOUR PROCEDURE:**

If you are taking any diet pills such as: Fenfluramine, Dexfenfluramine or Phentermine, stop taking those **2 weeks prior** to your colonoscopy. Failure to stop these medications may result in having to reschedule your procedure the day of.

***STOP THESE MEDICATIONS 7 DAYS BEFORE YOUR PROCEDURE:**

If you take GLP-1 Meds- Ozempic, Wegovy, Rybelsus, Trulicity, Byetta, Bydureon BCise, Saxenda, Victoza, Mounjaro or any other weekly injectables for weight loss / diabetes, you must contact your prescriber to stop this medication 7 days before your procedure. You may resume the medication after the procedure. If you have not stopped these medications 7 days prior, your procedure **WILL** be canceled per anesthesia. Daily diabetic medication instructions are under 1 day before/day of procedure section.

***STOP THESE MEDICATIONS 5 DAYS BEFORE YOUR PROCEDURE:**

Excedrin, Ibuprofen, Advil, Motrin, Aleve, Naprosyn, Naproxen, Celebrex, Celecoxib, Mobic, Meloxicam, Relafen, Nabumetone. Stop all herbal medications, supplements, and vitamins 5 days prior to your procedure. ****Tylenol is okay to take.**

Blood thinning medications including, Coumadin, Warfarin, Plavix, Xarelto, Pradaxa, Effient, Aggrenox, Eliquis or Brilinta must be stopped prior to your procedure as directed by the prescribing physician. Our office will send a request, but it is ultimately the **patient's responsibility** to also know if they can be off the blood thinner or not.

*****You may continue to take a low dose (81mg) Aspirin.**

***STOP SGLT-2 INHIBITOR MEDICATIONS 4 DAYS PRIOR TO YOUR PROCEDURE:**

Invokana (Canagliflozin), Farxiga (Dapagliflozin), Jardiance (Empagliflozin), Steglatro (ertugliflozin), Brenzavvy (Bexaglifoxin) Failure to stop these medications **WILL** result in a cancelled procedure per anesthesia.

FOODS TO AVOID 4 DAYS BEFORE YOUR PROCEDURE:

Fresh fruits with seeds, raw vegetables, nuts, popcorn, seeds, whole grain bread, whole grain pasta, lentils, beans and corn. (You may have apples, pears or peaches that are peeled. **Do Not eat the skins.**)

PURCHASE THESE NONPRESCRIPTION ITEMS BEFORE YOUR COLONOSCOPY:

- Dulcolax: 2 tablets (5mg Each) **Laxative** (not the stool softener)

1 DAY BEFORE YOUR COLONOSCOPY:

- If you are **diabetic and on oral medication (Metformin, Starlix, Prandin, Glipizide, Tradjenta, Januvia, Nesina, Avandia, Oseni, etc):** Take only half the usual dose the day BEFORE the procedure. **Do not take** your diabetic medication on the day of your procedure. You may take your medication when the procedure is over and you are able to eat.
- ✓ You will be on a limited diet the entire day. Allowed dairy and liquids listed below.
- ✓ Allowed dairy products: Eggs, milk (including soy, rice or almond), plain cheese (no cottage cheese), vanilla ice cream, plain or vanilla yogurt (without fruit, nuts or granola)
- ✓ Allowed clear Liquids: mineral, bubbly or plain water, Gatorade, apple juice, white grape juice, ginger ale, clear broth or bouillon (without veggies, meats or rice), Jell-o, all carbonated and noncarbonated soft drinks, plain popsicles, coffee and tea. Avoid all liquids or popsicles with red or purple dyes.
- ✓ You must start to hydrate yourself, which means you must start to drink extra fluids.
- ✓ **DO NOT DRINK: tomato juice, orange juice, grapefruit juice, protein, or drinks with fiber in them. NO beer, wine or alcohol is allowed. Nothing with Red or Purple Dyes.**

AT 4 PM:

- ✓ Take (2) two Dulcolax tablets. (5mg Each) Laxative. After you finish this portion of the prep, you may NOT have any solids by mouth. You may have clear liquids only.

AT 6 PM:

- ✓ Drink the entire first bottle of the Clenpiq fluid. Follow with a minimum of (5) 8-ounce glasses of clear fluid at your own pace over the next 5 hours. You may drink more if you choose. The prep may take up to 4 hours to begin working.
- ✓ You **MUST** drink more solution in the morning even if your stool is clear.

DAY OF YOUR COLONOSCOPY:

- ✓ **Start drinking the prep solution 5 hours before your test(_____:_____).** You must be done drinking the prep and additional fluids in 1 hour. Drink the entire second bottle of the Clenpiq fluid. Follow the prep with (4) 8-ounce glasses of clear fluid within the next hour. You must finish drinking your prep **at least 4 hours** before your procedure time. **NOTHING BY MOUTH – NO FOOD OR DRINK OF ANY SORT AFTER FINISHING YOUR PREP!**
- **You may take any medications prescribed for your heart, blood pressure, asthma, seizure disorder or thyroid with a small sip of water 4 hours prior to your scheduled procedure time.**
- **If you are diabetic and on insulin: Test your blood sugar before you come for your colonoscopy. Take only half of your usual dose of insulin on the day of your procedure.**



After you finish this portion of the prep, **you may NOT have any liquids or solids by mouth. NO FOOD, FLUID, GUM, MINTS, OR CANDY IS ALLOWED UNTIL AFTER THE PROCEDURE.** You will be rescheduled if you do not follow these instructions.