



Good Shepherd
West Pavilion
(450 IL-22, Barrington, IL
60010)

Northwestern Mchenry
Main Entrance
(4201 Medical Centre Dr, McHenry, IL
60050)

Physician: SPANGLER Date: _____ Arrival Time: _____ Procedure Time: _____

LOWER ENDOSCOPIC ULTRASOUND (EUS) PREP:

A lower endoscopic ultrasound (EUS) is a procedure that allows the doctor to examine the lower digestive tract using sound waves to create visual pictures of the organs involved in the digestive process. The doctor is looking for any abnormalities previously identified on a colonoscopy or x-ray tests like CT scans and may take biopsies.

IT IS ESSENTIAL THAT...

1. You bring your ID, insurance card and a list of medications (including dosages) to your appointment.
2. You will not be allowed to drive yourself home after the procedure due to the sedation. Therefore, you will need an adult to drive you home. You may **NOT** use public transportation. **If you do not have a ride, your procedure will be cancelled.**

MEDICATIONS TO AVOID PRIOR TO PROCEDURE: Failure to stop medications may result in being rescheduled the day of your procedure after you are prepped. Please review medications below and call with any questions. *Daily diabetic medication instructions can be found in 1 day before/day of procedure section.*

***STOP 14 DAYS BEFORE YOUR PROCEDURE:**

If you are taking any diet pills such as: Fenfluramine, Dexfenfluramine or Phentermine, stop taking those **2 weeks prior** to your colonoscopy. Failure to stop these medications may result in having to reschedule your procedure the day of.

***STOP 7 DAYS BEFORE YOUR PROCEDURE:**

If you take GLP-1 Meds- Ozempic, Wegovy, Rybelsus, Trulicity, Byetta, Bydureon BCise, Saxenda, Victoza, Mounjaro or any other weekly injectables for weight loss / diabetes, you must contact your prescriber to stop this medication 7 days before your procedure. You may resume the medication after the procedure. If you have not stopped these medications 7 days prior, your procedure **WILL** be canceled per anesthesia. Daily diabetic medication instructions are under 1 day before/day of procedure section.

***STOP 5 DAYS BEFORE YOUR PROCEDURE:**

Excedrin, Ibuprofen, Advil, Motrin, Aleve, Naprosyn, Naproxen, Celebrex, Celecoxib, Mobic, Meloxicam, Relafen, Nabumetone. Stop all herbal medications, supplements, and vitamins 5 days prior to your procedure. ****Tylenol is okay to take.**

Blood thinning medications including, Coumadin, Warfarin, Plavix, Xarelto, Pradaxa, Effient, Aggrenox, Eliquis or Brilinta must be stopped prior to your procedure as directed by the prescribing physician. Our office will send a request, but it is ultimately the **patients responsibility** to also know if they are able to be off the blood thinner or not.

*****You may continue to take a low dose (81mg) Aspirin.**

MORE →

***STOP SGLT-2 INHIBITOR MEDICATIONS 4 DAYS PRIOR TO YOUR PROCEDURE:**

Invokana (Canagliflozin), Farxiga (Dapagliflozin), Jardiance (Empagliflozin), Steglatro (ertugliflozin), Brenzavvy (Bexaglifoxin). Failure to stop these medications **WILL** result in a rescheduled procedure per anesthesia.

PURCHASE THIS NONPRESCRIPTION ITEM 4 DAYS BEFORE YOUR LOWER EUS:

- ✓ Two (2) fleet enemas from your pharmacy

1 DAY BEFORE YOUR LOWER EUS:

- If you are **diabetic and on oral medication (Metformin, Starlix, Prandin, Glipizide, Tradjenta, Januvia, Nesina, Avandia, Oseni, etc)**: Take only half the usual dose the day BEFORE the procedure. **Do not take** your diabetic medication on the day of your procedure. You may take your medication when the procedure is over and you are able to eat.
 - ✓ You may eat your usual diet.
 - ✓ No alcohol, beer, or wine today or tomorrow.
 - ✓ **After midnight NOTHING should be taken by mouth. No food, fluid, gum, or candy is allowed until after your lower EUS.**

DAY OF YOU LOWER EUS:

- ✓ **No food, fluid, gum, or candy is allowed until after your procedure.**
- ✓ **You may take any medications prescribed for your heart, blood pressure, asthma, seizure disorder or thyroid with a small sip of water 4 hours prior to your scheduled procedure time.**
- ✓ **If you are diabetic and on insulin: Test your blood sugar before you come for your procedure. Take only half of your usual dose of insulin on the day of your procedure.**
- ✓ **3 hours before your procedure _____: _____, administer the first Fleet Enema. After having a bowel movement, administer the 2nd Fleet Enema.**



After you finish this portion of the prep, **you may NOT have any liquids or solids by mouth. NO FOOD, FLUID, GUM, MINTS, OR CANDY IS ALLOWED UNTIL AFTER THE PROCEDURE.** You will be rescheduled if you do not follow these instructions.