

# GI PARTNERS

of ILLINOIS, LLC

Barrington Office  
Suite 312  
(22285 N Pepper Rd, Ste 312,  
Lake Barrington IL, 60010)

Good Shepherd  
West Pavilion  
(450 IL-22, Barrington, IL  
60010)

Northwestern Huntley  
Main Entrance  
(10400 Haligus Rd, Huntley, IL  
60142)

Algonquin Surgery Center  
Main Entrance  
(2550 W Algonquin Rd, Lake in the  
Hills IL, 60156)

Physician: \_\_\_\_\_ Date: \_\_\_\_\_ Arrival Time: \_\_\_\_\_ Procedure Time: \_\_\_\_\_

## **FLEXIBLE SIGMOIDOSCOPY - NO SEDATION PREP:**

A flexible sigmoidoscopy is a procedure that examines the rectum and the last third section of the large intestine called the sigmoid colon with a flexible fiberoptic instrument. The doctor is looking for any abnormalities in the lining of the sigmoid colon and rectum and may take biopsies.

## **\*\*\*STOP 5 DAYS BEFORE YOUR PROCEDURE:**

Excedrin, Ibuprofen, Advil, Motrin, Aleve, Naprosyn, Naproxen, Celebrex, Celecoxib, Mobic, Meloxicam, Relafen, Nabumetone. Stop all herbal medications, supplements, and vitamins 5 days prior to your procedure. **\*\*Tylenol is okay to take.**

Blood thinning medications including, Coumadin, Warfarin, Plavix, Xarelto, Pradaxa, Effient, Aggrenox, Eliquis or Brilinta must be stopped prior to your procedure as directed by the prescribing physician. Our office will send a request, but it is ultimately the **patients responsibility** to also know if they are able to be off the blood thinner or not.

**\*\*\*You may continue to take a low dose (81mg) Aspirin.**

## **PURCHASE THIS NONPRESCRIPTION ITEM 4 DAYS BEFORE YOUR FLEXI:**

- ✓ Two (2) fleet enemas from your pharmacy

## **1 DAY BEFORE YOUR FLEXI:**

- ✓ You may eat your usual diet.
- ✓ No alcohol, beer, or wine today or tomorrow.
- ✓ **After midnight NOTHING should be taken by mouth. No food, fluid, gum, or candy is allowed until after your Flexi.**

## **DAY OF YOUR FLEXI:**

- ✓ **3 hours before your procedure \_\_\_\_\_:\_\_\_\_\_**, administer the first Fleet Enema. After having a bowel movement, administer the 2nd Fleet Enema.
- ✓ **You may NOT have any liquids OR solids by mouth. NO FOOD, FLUID, GUM OR CANDY IS ALLOWED.**

**\*\*\*You may take any medications prescribed for your heart, blood pressure, asthma, seizure disorder or thyroid with a small sip of water 4 hours prior to your scheduled procedure time.**